INDIAN SCHOOL AL WADI AL KABIR
Assessment - 1

Class: XII | Sub: MATHEMATICS (041) | Max Marks: 80
Date: 30.08.2026 | (Sample Paper) | Time: 3 hr

General Instructions:

1. This question paper is divided in to 6 sections- A, B, C, D and E
2. Section A comprises of 20 MCQ type questions of 1 mark each.
3. Section B comprises of 5 Very Short Answer Type Questions of 2 marks each.
4. Section C comprises of 6 Short Answer Type Questions of 3 marks each.
5. Section D comprises of 4 Long Answer Type Questions of 5 marks each.
6. Section E comprises of 3 source based / case based / passage-based questions (4 marks each) with sub parts.
7. Internal choice has been provided for certain questions

SECTION – A
(Each MCQ Carries 1 Mark)

1 If $f'(x) = x + \frac{1}{x}$, then $f(x)$ is

a) $x^2 + \log|x| + c$ b) $\frac{x^2}{2} + \log|x| + c$ c) $\frac{x}{2} + \log|x| + c$ d) $\frac{x}{2} - \log|x| + c$

2 The value of 'k' for which the function $f(x) = \begin{cases} 3x + 5, & if\ x \geq 2 \\ kx^2, & if\ x < 2 \end{cases}$ is continuous is

a) $\frac{-11}{4}$ b) $\frac{11}{4}$ c) $\frac{-4}{11}$ d) $\frac{4}{11}$

3 Find k, if A = $\begin{bmatrix} -2 & 3 \\ k & 4 \end{bmatrix}$ is a singular matrix

a) $\frac{3}{8}$ b) $\frac{-3}{8}$ c) $\frac{-8}{3}$ d) $\frac{8}{3}$

4 The value of $\int_2^3 \frac{x}{x^2+1}\,dx$ is

a) log 4 b) log $\frac{3}{2}$ c) $\frac{1}{2}$ log 2 d) log $\frac{9}{4}$

5 The value of $\sin^{-1}\left[\sin\left(\frac{13\pi}{7}\right)\right]$ is

a) $\frac{13\pi}{7}$ b) $-\frac{13\pi}{7}$ c) $\frac{\pi}{7}$ d) $-\frac{\pi}{7}$

6 Find the maximum profit that a company can make, if the profit function is given by $P(x) = 72 + 42x - x^2$, where x is the number of units and P is the profit in rupees.

a) ₹ 42 b) ₹ 72 c) ₹ 114 d) ₹ 513

7 $\int \frac{(x-1)}{(x-2)(x-3)}$ dx equals

a) $2\log|x-3| - \log|x-2| + C$ c) $\log|x-3| - 2\log|x-2| + C$

b) $\log|x-3| - \log|x-2| + C$ d) $\log|x-2| - \log|x-3| + C$

8 If $[x-2 \quad 5+y]\begin{bmatrix}0 & 1\\ 1 & 0\end{bmatrix} = 0$, then x + y is

a) 0 b) -1 c) -2 d) -3

9 Let $\sin^{-1}(2x) + \cos^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{2}$. Then the value of *'x'* is

a) $\frac{1}{2}$ b) $\frac{1}{4}$ c) $\frac{1}{8}$ d) 8

10 If A is a square matrix of order 2 and |A| = - 3, then the value of |5A| is

a) -3 b) -15 c) -75 d) None of these

11 If $y = a\cos mx + b\sin mx$, then $\frac{d^2y}{dx^2}$ is

a) m^2y b) $-m^2y$ c) my d) - my

12 Value of $\int \frac{\cos\sqrt{x}}{\sqrt{x}}$ dx is

a) $-2\sin\sqrt{x} + c$ b) $\sin\sqrt{x} + c$ c) $2\cos\sqrt{x} + c$ d) $2\sin\sqrt{x} + c$

13 For the function $y = \frac{(x-7)}{(x-2)(x-3)}$, the value of $\left(\frac{dy}{dx}\right)_{\text{at } x = 7}$ is

a) - 20 b) $\frac{-1}{20}$ c) $\frac{1}{20}$ d) 20

14 The radius of the circle is increasing at the rate of 0.7 cm/sec. What is the rate of increase of its circumference?

a) 0.7 b) π c) 1.4π d) 2π

15 Find ***x*** if $\begin{vmatrix}3 & -6\\ 4 & 0\end{vmatrix} = \begin{vmatrix}3 & x^2\\ x & -1\end{vmatrix}$

a) 4 b) $\sqrt{-6}$ c) -3 d) None of these

16 The total revenue in rupees received from the sale of x units of a product is given by $R(x) = 13x^2 + 26x + 15$. Find the marginal revenue when $x = 7$.

a) 208 b) 54 c) 52 d) 13

17 If f(x) = log x, then $f^{l}(x) + f^{l}\left(\frac{1}{x}\right)$ is

a) $\frac{x^2-1}{x}$ a) $\frac{1-x^2}{x}$ a) $\frac{x^2+1}{x}$ a) $\frac{1+x}{x}$

18 If y = log $\sqrt{\tan x}$, then the value of $\frac{dy}{dx}$ at x = $\frac{\pi}{4}$

a) ∞ b) 1 c) $\frac{1}{2}$ d) 0

Directions: In the following 2 questions, A statement of Assertion (A) is followed by a statement of Reason (R). Mark the correct choice as.
(A) Both A and R are true and R is the correct explanation of A
(B) Both A and R are true but R is NOT the correct explanation of A
(C) A is true but R is false
(D) A is false and R is True

19 **Assertion (A):** If A is a square matrix such that $A^2 = A$, then $(I + A)^2 - 3A = I$
Reason (R): AI = IA = A

a) b) c) d)

20 **Assertion (A):** The relation f: {1, 2, 3, 4} → {x, y, z, p} defined by
f = {(1, x), (2, y), (3, z)} is a bijective function.
Reason (R): The relation f: {1, 2, 3} → {x, y, z, p} such that
f = {(1, x), (2, y), (3, z)} is one-one

a) b) c) d)

SECTION – B

(Each Question Carries 2 Marks)

21 If A = $\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & x \\ -2 & 2 & -1 \end{bmatrix}$ is a matrix satisfying AA' = 9I, find x

22 Evaluate $\int_0^1 \frac{x\, e^x}{(1+x)^2}$ dx

- OR -

Evaluate $\int_1^3 \frac{\sqrt[3]{x}}{\sqrt[3]{x}+\sqrt[3]{4-x}}$ dx

23 Find the value of $sin^{-1}\left[\cos\left(\frac{33\pi}{5}\right)\right]$

- OR -

Find the domain of $\sin^{-1}(x^2 - 4)$

24 Prove that the function *f* is surjective, where $f: N \to N$ such that

$$f(x) = \begin{cases} \frac{n+1}{2}, & if\ n\ is\ odd \\ \frac{n}{2}, & if\ n\ is\ even \end{cases}$$

Is the function injective? Justify your answer.

25 Find the value of '*a*' and '*b*' such that the function defined is a continuous function

$$f(x) = \begin{cases} 5, & if\ x \le 2 \\ ax + b, & if\ 2 < x < 10 \\ 21, & if\ x \ge 10 \end{cases}$$

SECTION – C

(Each Question Carries 3 Marks)

26 Find $\frac{dy}{dx}$ if $x = a\left(\cos t + \log\tan\frac{t}{2}\right)$ and $y = a \sin t$

27 Integrate the function $\int \frac{x^2+1}{x^2-5x+6}\,dx$

- OR -

Evaluate $\int_1^3 |x^3 - 2x|\,dx$

28 The volume of a cube is increasing at a rate of 9 cubic centimetres per second. How fast is the surface area increasing when the length of an edge is 10 centimetres?

- OR -

A particle moves along the curve $y = \frac{2}{3}x^3 + 1$. Find the coordinates of the points on the curve at which the y-coordinate is changing twice as fast as the x coordinate.

29 Let $f: N \to R$ be a function defined as $f(x) = 4x^2 + 12x + 15$, where R is the range of *f*. Show that the function *f(x)* is one-one.

- OR -

Check whether a function $f: R \to \left[-\frac{1}{2}, \frac{1}{2}\right]$ defined as $f(x) = \frac{x}{(1+x)^2}$ is one-one or onto or not

30 Express the matrix $A = \begin{bmatrix} 2 & 4 & -6 \\ 7 & 3 & 5 \\ 1 & -2 & 4 \end{bmatrix}$ as the sum of a symmetric and skew symmetric matrix.

31 If $y = 3\cos(\log x) + 4\sin(\log x)$, show that $x^2 y_2 + xy_1 + y = 0$

SECTION – D

(Each Question Carries 5 Marks)

32 Define the relation R in the set $N \times N$ as follows:
For (a, b), (c, d) $\in N \times N$, (a, b) R (c, d) iff ad = bc.
Prove that R is an equivalence relation in $N \times N$.

- OR -

Show that the function f: R $\rightarrow$ {x $\in$ R : - 1< x < 1} defined by f(x) = $\frac{x}{1+|x|}$,
x $\in$ R is a one-one onto function

33 The equilibrium conditions for three competitive markets are described as given below, where m_1, m_2 and m_3 are the equilibrium price for each market respectively.
$m_1 + 2m_2 + 3m_3 = 85$
$3m_1 + 2m_2 + 2m_3 = 105$
$2m_1 + 3m_2 + 2m_3 = 110$
Using matrix method, find the values of respective equilibrium prices

34 Find $\frac{dy}{dx}$ of the function $x^y + y^x = 1$

- OR -

If $y = Ae^{mx} + Be^{nx}$, show that $\frac{d^2y}{dx^2} - (m + n)\frac{dy}{dx} + mny = 0$

35 Integrate the function $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}}\, dx$

SECTION – E

(CASE STUDY - Each Question Carries 4 Marks)

36 Read the following passage and answer the questions given below.

The temperature of a person during an intestinal illness is given by
$f(x) = -0.1x^2 + mx + 98.6$, $0 \le x \le 12$, m being a constant, where f(x) is the temperature in F^0 at x days.

(i) Is the function differentiable in the interval (0, 12)? Justify your answer (1m)
(ii) If 6 is the critical point of the function, then find the value of the constant m (1m)
(iii) Find the intervals in which the function is strictly increasing / strictly decreasing. (2m)

- OR -

Find the points of local maximum / local minimum, if any, in the interval (0, 12) as well as the points of absolute maximum / absolute minimum in the interval [0, 12]. Also, find the corresponding local maximum / local minimum. (2m)

37 For an EMC project, a student of Class XII makes an open cardboard box for a jewelry shop from a square sheet of side 18 cm by cutting off squares from each corner and folding up the flaps. Assume that 'x' be the side of squares cut off from each corner. Based on the given information, answer the following questions.



(i) For the open box, find the length, breadth and height in terms of x. (1m)

(ii) Write an expression for the volume of the open box. (1m)

(iii) For what value of 'x', the open box will have maximum volume? (2m)

- OR -

Find the maximum value of volume of the open box. (2m)

38 Gautam buys 5 pens, 3 bags and 1 instrument box and pays a sum of ₹160. From the same shop, Vikram buys 2 pens, 1 bag and 3 instrument boxes and pays a sum of ₹190. Also, Ankur buys 1 pen, 2 bags and 4 instrument boxes and pays a sum of ₹250. Based on the above information, answer the following questions.

(i) Convert the given above situation into a matrix equation of the form AX = B (1m)

(ii) Find |A| (1m)

(iii) Find A^{-1} (2m)

- OR -

Determine $P = A^2 - 5A$ (2m)

INDIAN SCHOOL AL WADI AL KABIR

Assessment - 1

Class: XII | Sub: MATHEMATICS (041) | Max Marks: 80

Date: 30.08.2026 | (Answers) | Time: 3 hr

1	b) $\frac{x^2}{2}$ + log |x| + c
2	b) $\frac{11}{4}$
3	c) $\frac{-8}{3}$
4	c) $\frac{1}{2}\log 2$
5	d) $-\frac{\pi}{7}$
6	d) ₹ 513
7	a) $2\log\|x-3\| - \log\|x-2\| + C$
8	d) -3
9	b) $\frac{1}{4}$
10	c) -75
11	b) $-m^2y$
12	d) 2 sin √x + c
13	c) $\frac{1}{20}$
14	d) 2π
15	c) -3
16	a) 208
17	a) $\frac{x^2+1}{x}$
18	b) 1
19	(A) Both A and R are true and R is the correct explanation of A
20	(D) A is false and R is True
21	$AA' = \begin{bmatrix}1&2&2\\2&1&x\\-2&2&-1\end{bmatrix}\begin{bmatrix}1&2&-2\\2&1&2\\2&x&-1\end{bmatrix} = 9\begin{bmatrix}1&0&0\\0&1&0\\0&0&1\end{bmatrix}$ $\begin{bmatrix}9&2x+4&0\\2x+4&x^2+5&-x-2\\0&-x-2&9\end{bmatrix} = \begin{bmatrix}9&0&0\\0&9&0\\0&0&9\end{bmatrix}$ $2x+4=0$ $x=-2$
22	$I = \int_0^1 \frac{xe^x}{(1+x)^2}dx$ $= \int_0^1 \frac{(x+1)-1}{(1+x)^2}e^x dx$ $= \int_0^1 \left[\frac{(x+1)e^x}{(x+1)^2} - \frac{e^x}{(x+1)^2}\right]dx$ $= \int_0^1 e^x\left[\frac{1}{1+x} - \frac{1}{(1+x)^2}\right]dx$: Let $I = \int_1^3 \frac{\sqrt[3]{x}}{\sqrt[3]{x}+\sqrt[3]{4-x}}dx$ .....(1) $I = \int_1^3 \frac{\sqrt[3]{4-x}}{\sqrt[3]{4-x}+\sqrt[3]{4-(4-x)}}dx$ $I = \int_1^3 \frac{\sqrt[3]{4-x}}{\sqrt[3]{4-x}+\sqrt[3]{x}}dx$ .....(2)

$$= \left[e^x \frac{1}{1+x} \right]_0^1$$

$$\left[\because \int e^x [f(x) + f'(x)]\,dx = e^x f(x) + c \right]$$

$$= \frac{e^1}{1+1} - \frac{e^0}{1+0} = \frac{e}{2} - 1$$

$$I = \frac{e}{2} - 1$$

Adding (1) and (2), we get

$$2I = \int_1^3 \frac{\sqrt[3]{x} + \sqrt[3]{4-x}}{\sqrt[3]{4-x} + \sqrt[3]{x}}\,dx$$

$$2I = \int_1^3 1\,dx$$

$$2I = x|_1^3$$

$2I = 3 - 1 = 2$ gives $I = 1$

23

$$\sin^{-1}\left(\cos\left(\frac{33\pi}{5}\right)\right) = \sin^{-1}\cos\left(6\pi + \frac{3\pi}{5}\right) = \sin^{-1}\cos\left(\frac{3\pi}{5}\right) = \sin^{-1}\sin\left(\frac{\pi}{2} - \frac{3\pi}{5}\right)$$

$$= \frac{\pi}{2} - \frac{3\pi}{5} = -\frac{\pi}{10}.$$

OR

$$-1 \le (x^2 - 4) \le 1 \Rightarrow 3 \le x^2 \le 5 \Rightarrow \sqrt{3} \le |x| \le \sqrt{5}$$

$\Rightarrow x \in \left[-\sqrt{5}, -\sqrt{3}\right] \cup \left[\sqrt{3}, \sqrt{5}\right]$. So, required domain is $\left[-\sqrt{5}, -\sqrt{3}\right] \cup \left[\sqrt{3}, \sqrt{5}\right]$.

24

Let $y \in N$(codomain). Then $\exists\ 2y \in N$(domain) such that $f(2y) = \frac{2y}{2} = y$. Hence, f is surjective.

$1, 2 \in N$(domain) such that $f(1) = 1 = f(2)$

Hence, f is not injective.

25

Given that the function is continuous at $x = 2$.

Therefore, LHL = RHL $= f(2)$

$$\Rightarrow \lim_{x \to 2^-} f(x) = \lim_{x \to 2^+} f(x) = f(2)$$

$$\Rightarrow \lim_{x \to 2^-} 5 = \lim_{x \to 2^+} ax + b = 5$$

$$\Rightarrow 2a + b = 5 \quad \text{...(i)}$$

Given that the function is continuous at $x = 10$.

Therefore, LHL = RHL $= f(10)$

$$\Rightarrow \lim_{x \to 10^-} f(x) = \lim_{x \to 10^+} f(x) = f(10)$$

$$\Rightarrow \lim_{x \to 10^-} ax + b = \lim_{x \to 10^+} 21 = 21$$

$$\Rightarrow 10a + b = 21 \quad \text{...(ii)}$$

Solving the equation (i) and (ii), we get $a = 2$ $b = 1$

26

here, $x = a\left(\cos t + \log \tan \frac{t}{2}\right)$

Therefore, $\frac{dx}{dt} = a\left(-\sin t + \frac{1}{\tan \frac{t}{2}} . \sec^2 \frac{t}{2} . \frac{1}{2}\right) = a\left(-\sin t + \frac{\cos \frac{t}{2}}{\sin \frac{t}{2}} . \frac{1}{\cos^2 \frac{t}{2}} . \frac{1}{2}\right)$

$$= a\left(-\sin t + \frac{1}{2 \sin \frac{t}{2} \cos \frac{t}{2}}\right) \qquad = a\left(-\sin t + \frac{1}{\sin t}\right)$$

$$= a\left(\frac{-\sin^2 t + 1}{\sin t}\right) \qquad = a\left(\frac{\cos^2 t}{\sin t}\right)$$

$y = a \sin t \qquad \frac{dy}{dt} = a \cos t \qquad \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{a \cos t}{a\left(\frac{\cos^2 t}{\sin t}\right)} = \frac{\sin t}{\cos t} = \tan t$

27

$$\frac{x^2+1}{x^2-5x+6} = 1+\frac{5x-5}{x^2-5x+6}$$

$$=1+\frac{5x-5}{(x-2)(x-3)}$$

$$\frac{5x-5}{(x-2)(x-3)} = \frac{A}{x-2}+\frac{B}{x-3}$$

$5x-5 = A(x-3)+B(x-2)$

Equating the coefficients of x $A+B=5$

Equating the constant terms $3A+2B=5$

Solving $A=-5$ and $B=10$

$$\int\frac{x^2+1}{x^2-5x+6}dx = \int dx-5\int\frac{1}{x-2}dx+10\int\frac{dx}{x-3}$$

$= x-5\log|x-2|+10\log|x-3|+C.$

-OR-

Consider $I=\int_1^3|x^2-2x|\,dx$

$$|x^2-2x| = \begin{cases}-(x^2-2x) & \text{where } 1\le x<2\\ (x^2-2x) & \text{where } 2\le x\le 3\end{cases}$$

$$I=\int_1^2|x^2-2x|\,dx+\int_2^3|x^2-2x|\,dx$$

$$I=\int_1^2-(x^2-2x)\,dx+\int_2^3(x^2-2x)\,dx$$

$$I=-\left[\frac{x^3}{3}-x^2\right]_1^2+\left[\frac{x^3}{3}-x^2\right]_2^3$$

$$I=-\left[\frac{8}{3}-4-\frac{1}{3}+1\right]+\left[9-9-\frac{8}{3}+4\right]$$

$$=-\frac{7}{3}+3-\frac{8}{3}+4$$

$I=7-5=2$

28

Let x be the length of a side, V be the volume and S be the surface area of the cube.
$V = x^3$ and $S = 6x^2$, where x is a function of time t.

$\frac{dV}{dt} = 9\text{cm}^3/\text{s}$ (Given)

$$9=\frac{dV}{dt}=\frac{d}{dt}(x^3) = \frac{d}{dx}(x^3)\cdot\frac{dx}{dt}$$

$$=3x^2\cdot\frac{dx}{dt}$$

$$\frac{dx}{dt}=\frac{3}{x^2}$$

$$\frac{dS}{dt}=\frac{d}{dt}(6x^2) = \frac{d}{dx}(6x^2)\cdot\frac{dx}{dt}$$

$$=12x\cdot\left(\frac{3}{x^2}\right)=\frac{36}{x}$$

when $x = 10$ cm,

$$\frac{dS}{dt}=3.6\,\text{cm}^2/\text{s}$$

- OR -

We have $y=\frac{2}{3}x^3+1$ ...(i)

$$\Rightarrow\frac{dy}{dt}=\frac{2}{3}\times3x^2\times\frac{dx}{dt}=2x^2\frac{dx}{dt}$$

$$\because\frac{dy}{dt}=2\frac{dx}{dt}$$

$$\Rightarrow 2\frac{dx}{dt}=2x^2\frac{dx}{dt}$$

$\Rightarrow x^2=1 \qquad \therefore x=\pm1$

Substituting values of x in (i),

we get : $y=\frac{5}{3},\frac{1}{3}$

points on the curve are $\left(1,\frac{5}{3}\right)$ and $\left(-1,\frac{1}{3}\right)$

29

Here $f:N\to R$, $f(x)=4x^2+12x+15$

Let $x_1,x_2\in N$ and $f(x_1)=f(x_2)$

Then, $4x_1^2+12x_1+15=4x_2^2+12x_2+15$

$\Rightarrow 4x_1^2+12x_1=4x_2^2+12x_2$

$\Rightarrow 4(x_1^2-x_2^2)+12(x_1-x_2)=0$

$\Rightarrow (x_1-x_2)[4(x_1+x_2)+12]=0$

As $x_1,x_2\in N$ so, $4(x_1+x_2)+12\neq0$

$\therefore (x_1-x_2)=0$

$\Rightarrow x_1=x_2$ so, f is one-one.

$$f(2)=\frac{2}{5},\ f\left(\frac{1}{2}\right)=\frac{\frac{1}{2}}{1+\left(\frac{1}{2}\right)^2}=\frac{1/2}{5/4}=\frac{2}{5}$$

That is, $f(2)=f\left(\frac{1}{2}\right)$ but $2\neq\frac{1}{2}$.

Hence, f is not one-one.

Now let $y=f(x)$, $y\in\left[-\frac{1}{2},\frac{1}{2}\right]$

That is, $y=\frac{x}{1+x^2}$

	-OR -	For this quadratic equation in x, for all $x \in R$ we must have $(-1)^2 - 4y \times y \geq 0$ That is, $1 - 4y^2 \geq 0$ $\Rightarrow (1-2y)(1+2y) \geq 0$ That is, $y \in \left[-\frac{1}{2}, \frac{1}{2}\right]$. That means, Range = Codomain . Hence, f is onto.
30	$A = \begin{bmatrix} 2 & 4 & -6 \\ 7 & 3 & 5 \\ 1 & -2 & 4 \end{bmatrix}$ $A' = \begin{bmatrix} 2 & 7 & 1 \\ 4 & 3 & -2 \\ -6 & 5 & 4 \end{bmatrix}$ $P = \frac{1}{2}(A+A') = \frac{1}{2}\begin{bmatrix} 4 & 11 & -5 \\ 11 & 6 & 3 \\ -5 & 3 & 8 \end{bmatrix}$ $= \begin{bmatrix} 2 & \frac{11}{2} & -\frac{5}{2} \\ \frac{11}{2} & 3 & \frac{3}{2} \\ -\frac{5}{2} & \frac{3}{2} & 4 \end{bmatrix}$ Since $P' = P$ $\therefore P$ is a symmetric matrix.	Let $Q = \frac{1}{2}(A-A') = \frac{1}{2}\begin{bmatrix} 0 & -3 & -7 \\ 3 & 0 & 7 \\ 7 & -7 & 0 \end{bmatrix}$ $= \begin{bmatrix} 0 & -\frac{3}{2} & -\frac{7}{2} \\ \frac{3}{2} & 0 & \frac{7}{2} \\ \frac{7}{2} & -\frac{7}{2} & 0 \end{bmatrix}$ $Q' = -\begin{bmatrix} 0 & -\frac{3}{2} & -\frac{7}{2} \\ \frac{3}{2} & 0 & \frac{7}{2} \\ \frac{7}{2} & -\frac{7}{2} & 0 \end{bmatrix} = -Q$ Since $Q' = -Q$ $\therefore Q$ is a skew symmetric matrix.
	Also $P+Q = \begin{bmatrix} 2 & \frac{11}{2} & -\frac{5}{2} \\ \frac{11}{2} & 3 & \frac{3}{2} \\ -\frac{5}{2} & \frac{3}{2} & 4 \end{bmatrix} + \begin{bmatrix} 0 & -\frac{3}{2} & -\frac{7}{2} \\ \frac{3}{2} & 0 & \frac{7}{2} \\ \frac{7}{2} & -\frac{7}{2} & 0 \end{bmatrix}$ $= \begin{bmatrix} 2 & 4 & -6 \\ 7 & 3 & 5 \\ 1 & -2 & 4 \end{bmatrix} = A$	
31	Given that $y = 3\cos(\log x) + 4\sin(\log x)$ $\frac{dy}{dx} = \frac{d}{dx}(3\cos(\log x) + 4\sin(\log x)) = -3\sin(\log x).\frac{1}{x} + 4\cos(\log x).\frac{1}{x}$ $\Rightarrow x\frac{dy}{dx} = -3\sin(\log x) + 4\cos(\log x)$ $x\frac{d^2y}{dx^2} + \frac{dy}{dx}.\frac{d}{dx}x = \frac{d}{dx}[-3\sin(\log x) + 4\cos(\log x)]$ $= -3\cos(\log x).\frac{1}{x} - 4\sin(\log x).\frac{1}{x} = -\frac{1}{x}[3\cos(\log x) + 4\sin(\log x)] = -\frac{1}{x}.y$ $\Rightarrow x\frac{d^2y}{dx^2} + \frac{dy}{dx} = -\frac{1}{x}y$ $\Rightarrow x^2\frac{d^2y}{dx^2} + x\frac{dy}{dx} = -y$ $\Rightarrow x^2y_2 + xy_1 + y = 0$	

32

Let $(a, b) \in N \times N$. Then we have

ab = ba (by commutative property of multiplication of natural numbers)

$\Rightarrow (a, b)R(a, b)$

Hence, R is reflexive.

Let $(a, b), (c, d) \in N \times N$ such that (a, b) R (c, d). Then

ad = bc

$\Rightarrow cb = da$ (by commutative property of multiplication of natural numbers

$\Rightarrow (c, d)R(a, b)$

Hence, R is symmetric.

Let $(a, b), (c, d), (e, f) \in N \times N$ such that (a, b) R (c, d) and (c, d) R (e, f).

Then ad = bc, cf = de

$\Rightarrow adcf = bcde$

$\Rightarrow af = be$

$\Rightarrow (a, b)R(e, f)$

Hence, R is transitive.

Since, R is reflexive, symmetric and transitive, R is an equivalence relation on $N \times N$.

- OR -

$|x| = \begin{cases} x & ,x \geq 0 \\ -x & ,x < 0 \end{cases} \Rightarrow f(x) = \begin{cases} \frac{x}{1+x}, & x \geq 0 \\ \frac{x}{1-x}, & x < 0 \end{cases}$

For x ≥ 0

$f(x_1) = \frac{x_1}{1+x_1}$

$f(x_2) = \frac{x_2}{1+x_2}$

$f(x_1) = f(x_2)$

$\frac{x_1}{1+x_1} = \frac{x_2}{1+x_2}$

$x_1(1 + x_2) = x_2(1 + x_1)$

$x_1 + x_1x_2 = x_2 + x_2x_1$

$x_1 = x_2$

For x < 0

$f(x_1) = \frac{x_1}{1-x_1}$

$f(x_2) = \frac{x_2}{1-x_2}$

Putting $f(x_1) = f(x_2)$

$\frac{x_1}{1-x_1} = \frac{x_2}{1-x_2}$

$x_1(1 - x_2) = x_2(1 - x_1)$

$x_1 - x_1x_2 = x_2 - x_2x_1$

$x_1 = x_2$

Hence, if $f(x_1) = f(x_2)$, then $x_1 = x_2$ ∴ f is **one-one**

For x ≥ 0

$f(x) = \frac{x}{1+x}$

Let f(x) = y,

$y = \frac{x}{1+x}$

$x = \frac{y}{1-y}$, for x ≥ 0

For x < 0

$f(x) = \frac{x}{1-x}$

Let f(x) = y

$y = \frac{x}{1-x}$

$x = \frac{y}{1+y}$, for x < 0

Here, $y \in \{x \in \mathbf{R}: -1 < x < 1\}$

So, x is defined for all values of y.

∴ f is **onto**

Hence, f is one-one and onto.

33

$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 2 \\ 2 & 3 & 2 \end{bmatrix}$

$\Rightarrow |A| = 9 \Rightarrow A^{-1}$ exists

And $A^{-1} = \frac{1}{9}\begin{bmatrix} -2 & 5 & -2 \\ -2 & -4 & 7 \\ 5 & 1 & -4 \end{bmatrix}$

$AX = B \Rightarrow X = A^{-1}B$

$\Rightarrow X = \frac{1}{9}\begin{bmatrix} -2 & 5 & -2 \\ -2 & -4 & 7 \\ 5 & 1 & -4 \end{bmatrix}\begin{bmatrix} 85 \\ 105 \\ 110 \end{bmatrix} = \begin{bmatrix} 15 \\ 20 \\ 10 \end{bmatrix}$

$\Rightarrow p_1 = 15, p_2 = 20, p_3 = 10$

34

Let $u = x^y$ and $v = y^x$, therefore, $u + v = 1$

Differentiating with respect to x, we get $\frac{du}{dx} + \frac{dv}{dx} = 0$ ...(i)

$u = x^y$, taking log on both the sides, $\log u = y \log x$,

$\frac{1}{u}\frac{du}{dx} = y.\frac{d}{dx}\log x + \log x.\frac{d}{dx}y = y.\frac{1}{x} + \log x.\frac{dy}{dx}$

$\frac{du}{dx} = x^y\left[\frac{y}{x} + \log x.\frac{dy}{dx}\right]$ ...(ii)

$v = y^x$, taking log on both the sides $\log v = x \log y$,

$\frac{1}{v}\frac{dv}{dx} = x.\frac{d}{dx}\log y + \log y.\frac{d}{dx}x = x.\frac{1}{y}\frac{dy}{dx} + \log y.1$

$\frac{dv}{dx} = v\left[\frac{x}{y}\frac{dy}{dx} + \log y\right] = y^x\left[\frac{x}{y}\frac{dy}{dx} + \log y\right]$ ...(iii)

Putting the value of $\frac{du}{dx}$ from (ii) and $\frac{dv}{dx}$ from (iii) in equation (i), we get

$x^y\left[\frac{y}{x} + \log x.\frac{dy}{dx}\right] + y^x\left[\frac{x}{y}\frac{dy}{dx} + \log y\right] = 0 \Rightarrow yx^{y-1} + x^y \log x\frac{dy}{dx} + xy^{x-1}\frac{dy}{dx} + y^x \log y = 0$

$\Rightarrow \frac{dy}{dx}(x^y \log x + xy^{x-1}) = -(y^x \log y + yx^{y-1}) \Rightarrow \frac{dy}{dx} = -\frac{y^x \log y + yx^{y-1}}{x^y \log x + xy^{x-1}}$

- OR -

$\frac{dy}{dx} = \frac{d}{dx}(Ae^{mx} + Be^{nx}) = mAe^{mx} + nBe^{nx}$

$\Rightarrow \frac{d^2y}{dx^2} = \frac{d}{dx}(mAe^{mx} + nBe^{nx}) = m^2Ae^{mx} + n^2Be^{nx}$

Putting the value of $\frac{d^2y}{dx^2}$ and $\frac{dy}{dx}$ in $\frac{d^2y}{dx^2} - (m+n)\frac{dy}{dx} + mny$

$\text{LHS} = (m^2Ae^{mx} + n^2Be^{nx}) - (m+n)(mAe^{mx} + nBe^{nx}) + mny$

$= m^2Ae^{mx} + n^2Be^{nx} - (m^2Ae^{mx} + mnBe^{nx} + mnAe^{mx} + n^2Be^{nx}) + mny$

$= -(mnAe^{mx} + mnBe^{nx}) + mny = -mn(Ae^{mx} + Be^{nx}) + mny$

$= -mny + mny = 0 = \text{RHS}$

35

Let $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$

$\Rightarrow I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{(\sin x + \cos x)}{\sqrt{-(-\sin 2x)}} dx$

$\Rightarrow I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{-(-1+1-2\sin x \cos x)}} dx$

$\Rightarrow I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{(\sin x + \cos x)}{\sqrt{1-(\sin^2 x + \cos^2 x - 2\sin x \cos x)}} dx$

$\Rightarrow I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{(\sin x + \cos x)dx}{\sqrt{1-(\sin x - \cos x)^2}}$

Let $(\sin x - \cos x) = t \Rightarrow (\sin x + \cos x)dx = dt$

When $x = \frac{\pi}{6}, t = \left(\frac{1-\sqrt{3}}{2}\right)$,

when $x = \frac{\pi}{3}, t = \left(\frac{\sqrt{3}-1}{2}\right)$

$$I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x+\cos x}{\sqrt{\sin 2x}}\,dx \quad = \int_{\frac{1-\sqrt{3}}{2}}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}}$$

$$I = \int_{-\left(\frac{\sqrt{3}-1}{2}\right)}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}}$$

As $\frac{1}{\sqrt{1-(-t)^2}} = \frac{1}{\sqrt{1-t^2}}$, hence, $\frac{1}{\sqrt{1-t^2}}$ is an even function

As we know that if $f(x)$ is an even function, then

$\int_{-a}^{a} f(x)dx = 2\int_{0}^{a} f(x)dx$

$$\Rightarrow I = 2\int_{0}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}} \quad = [2\sin^{-1} t]_0^{\frac{\sqrt{3}-1}{2}}$$

$$= 2\sin^{-1}\left(\frac{\sqrt{3}-1}{2}\right)$$

36

(i) $f(x) = -0.1x^2 + mx + 98.6$, being a polynomial function, is differentiable everywhere, hence, differentiable in (0, 12)

(ii) $f'(x) = -0.2x + m$

Since, 6 is the critical point,

$f'(6) = 0 \Rightarrow m = 1.2$

(iii) $f(x) = -0.1x^2 + 1.2x + 98.6$

$f'(x) = -0.2x + 1.2 = -0.2(x - 6)$

In the Interval	$f'(x)$	Conclusion
(0, 6)	+ve	f is strictly increasing in [0, 6]
(6, 12)	-ve	f is strictly decreasing in [6, 12]

OR

(iii) $f(x) = -0.1x^2 + 1.2x + 98.6$,

$f'(x) = -0.2x + 1.2, f'(6) = 0,$

$f''(x) = -0.2$

$f''(6) = -0.2 < 0$

Hence, by second derivative test 6 is a point of local maximum. The local maximum value = $f(6) = -0.1 \times 6^2 + 1.2 \times 6 + 98.6 = 102.2$

37	(i) For the open box the length, breadth and height is given by $(18-2x)$ cm, $(18-2x)$ cm and x cm respectively. (ii) Therefore, the volume of box is, $V=(18-2x)(18-2x)(x)=(324x-72x^2+4x^3)\ cm^3$ (iii) Now $\frac{dV}{dx}=324-144x+12x^2$ and $\frac{d^2V}{dx^2}=-144+24x$ For $\frac{dV}{dx}=0,\ 12(x^2-12x+27)=0$ $\Rightarrow (x-9)(x-3)=0$ Either $(x-9)=0$ or, $(x-3)=0$ $\because x\neq 9\ \ \therefore x=3$ cm $\because \left(\frac{d^2V}{dx^2}\right)_{at\ x=3}=-144+24(3)=-72<0$ So, V is maximum at $x=3$ cm. **OR** (iii) Refer the solution of (iii) as shown above. Clearly, the maximum volume of open box will be $V=(18-2x)(18-2x)(x)=(18-6)^2(3)$ $\Rightarrow V=432\ cm^3$.	
38	(i)	$\begin{bmatrix}5&3&1\\2&1&3\\1&2&4\end{bmatrix}\begin{bmatrix}x\\y\\z\end{bmatrix}=\begin{bmatrix}160\\190\\250\end{bmatrix}$
	(ii)	- 22
	(iii) – a	$A^{-1}=\begin{bmatrix}\frac{1}{11}&\frac{5}{11}&-\frac{4}{11}\\\frac{5}{22}&-\frac{19}{22}&\frac{13}{22}\\-\frac{3}{22}&\frac{7}{22}&\frac{1}{22}\end{bmatrix}$
	(ii) – b	$P=\begin{bmatrix}7&5&13\\5&8&2\\8&3&3\end{bmatrix}$